

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA700451

## CA12 Biotinylated Mouse Monoclonal Detection Antibody (Biotin conjugated) [Clone ID: OTI3F1]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI3F1
Applications:	ELISA
Recommended Dilution:	ELISA 1:1000
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CA12(NP_001209) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Predicted Protein Size:	39.5 kDa
Gene Name:	carbonic anhydrase 12
Database Link:	<u>NP 001209</u> <u>Entrez Gene 771 Human</u> <u>O43570</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CA12 Biotinylated Mouse Monoclonal Detection Antibody (Biotin conjugated) [Clone ID: OTI3F1] – TA700451
Background:	Carbonic anhydrase 12 (CA12) is an enzyme that in humans is encoded by the CA12 gene. Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. This gene product is a type I membrane protein that is highly expressed in normal tissues, such as kidney, colon and pancreas, and has been found to be overexpressed in 10% of clear cell renal carcinomas. Loss of function mutations in the CA12 gene result in defects in fluids and carbonate secretions, which is involved in cystic fibrosis-like syndrome with normal cystic fibrosis transmembrane conductance regulator (CFTR) protein levels, pancreatitis, Sjögren's syndrome and Xerostomia or dry mouth syndrome. It has been shown that CA12 expression is associated with a better prognosis in an unselected series of invasive breast carcinoma patients. Moreover, serum CA12 levels have been found significantly higher in lung cancer patients than in healthy controls, supporting that CA12 may be a promising diagnostic marker for lung cancer.
Synonyms:	CA-XII; CAXII; HsT18816; T18816
Protein Families	: Druggable Genome, Transmembrane
Protein Pathway	vs: Nitrogen metabolism

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US