

Product datasheet for TA800029M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

cIAP2 (BIRC3) Mouse Monoclonal Antibody [Clone ID: OTI1F5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1F5

Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human
Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 255-529 of human

BIRC3 (NP 001156) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 68.2 kDa

Gene Name: baculoviral IAP repeat containing 3

Database Link: NP 001156

Entrez Gene 330 Human

013489

cIAP2 (BIRC3) Mouse Monoclonal Antibody [Clone ID: OTI1F5] - TA800029M

Background: The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis

by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. The amino acid sequence predicts three baculovirus IAP repeat domains and a ring finger domain. Transcript variants encoding the

same isoform have been identified. [provided by RefSeq]


Synonyms: AIP1; API2; c-IAP2; CIAP2; HAIP1; HIAP1; MALT2; MIHC; RNF49

Protein Families: Druggable Genome

Protein Pathways: Apoptosis, Focal adhesion, NOD-like receptor signaling pathway, Pathways in cancer, Small

cell lung cancer, Ubiquitin mediated proteolysis

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BIRC3 ([RC207764], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BIRC3. Positive lysates [LY420093] (100ug) and [LC420093] (20ug) can be purchased separately from OriGene.