

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA800031S

cIAP2 (BIRC3) Mouse Monoclonal Antibody [Clone ID: OTI1D12]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI1D12	
Applications:	ons: IF, WB	
Recommended Dilution:	lution: WB 1:2000, IF 1:100	
Reactivity:	Human	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	luman recombinant protein fragment corresponding to amino acids 255-529 of human BIRC3 (NP_001156) produced in E.coli.	
Formulation:	oH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	tore at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	68.2 kDa	
Gene Name:	baculoviral IAP repeat containing 3	
Database Link:	Elink: <u>NP 001156</u> Entrez Gene 330 Human Q13489	




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	cIAP2 (BIRC3) Mouse Monoclonal Antibody [Clone ID: OTI1D12] – TA800031S
Background:	The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. The amino acid sequence predicts three baculovirus IAP repeat domains and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq]
Synonyms:	AIP1; API2; c-IAP2; CIAP2; HAIP1; HIAP1; MALT2; MIHC; RNF49
Protein Families:	Druggable Genome
Protein Pathway	s: Apoptosis, Focal adhesion, NOD-like receptor signaling pathway, Pathways in cancer, Small cell lung cancer, Ubiquitin mediated proteolysis

Product images:

170	_
130	_
100	_
70	-
55	-
40	-
35	-
25	-
15	-
10	_

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BIRC3 ([RC207764], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BIRC3. Positive lysates [LY420093] (100ug) and [LC420093] (20ug) can be purchased separately from OriGene.



Anti-BIRC3 mouse monoclonal antibody ([TA800031]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY BIRC3 ([RC207764]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US