

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA800039

### cIAP2 (BIRC3) Mouse Monoclonal Antibody [Clone ID: OTI2G7]

### **Product data:**

Product Type:	Primary Antibodies		
Clone Name:	OTI2G7		
Applications:	WB		
Recommended Dilution:	WB 1:2000		
Reactivity:	Human		
Host:	Mouse		
lsotype:	lgG1		
Clonality:	Monoclonal		
Immunogen:	Human recombinant protein fragment corresponding to amino acids 255-529 of human BIRC3 (NP_001156) produced in E.coli.		
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.		
Concentration:	1 mg/ml		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Conjugation:	Unconjugated		
Storage:	Store at -20°C as received.		
Stability:	Stable for 12 months from date of receipt.		
Predicted Protein Size:	68.2 kDa		
Gene Name:	baculoviral IAP repeat containing 3		
Database Link:	<u>NP_001156</u> <u>Entrez Gene 330 Human</u> <u>Q13489</u>		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	cIAP2 (BIRC3) Mouse Monoclonal Antibody [Clone ID: OTI2G7] – TA800039
Background:	The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. The amino acid sequence predicts three baculovirus IAP repeat domains and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq]
Synonyms:	AIP1; API2; c-IAP2; CIAP2; HAIP1; HIAP1; MALT2; MIHC; RNF49
Protein Families	: Druggable Genome
Protein Pathway	<b>s:</b> Apoptosis, Focal adhesion, NOD-like receptor signaling pathway, Pathways in cancer, Small cell lung cancer, Ubiquitin mediated proteolysis

## **Product images:**

170	-	
130	-	
100	-	
70	-	-
55	-	-
40	-	
35	-	
25	-	
15	-	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BIRC3 ([RC207764], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BIRC3. Positive lysates [LY420093] (100ug) and [LC420093] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US