

# **Product datasheet for TA800039BM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### cIAP2 (BIRC3) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2G7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2G7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 255-529 of human

BIRC3 (NP 001156) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 68.2 kDa

**Gene Name:** baculoviral IAP repeat containing 3

Database Link: NP 001156

Entrez Gene 330 Human

013489





**Background:** The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis

by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. The amino acid sequence predicts three baculovirus IAP repeat domains and a ring finger domain. Transcript variants encoding the

same isoform have been identified. [provided by RefSeq]


Synonyms: AIP1; API2; c-IAP2; CIAP2; HAIP1; HIAP1; MALT2; MIHC; RNF49

**Protein Families:** Druggable Genome

**Protein Pathways:** Apoptosis, Focal adhesion, NOD-like receptor signaling pathway, Pathways in cancer, Small

cell lung cancer, Ubiquitin mediated proteolysis

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BIRC3 ([RC207764], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BIRC3. Positive lysates [LY420093] (100ug) and [LC420093] (20ug) can be purchased separately from OriGene.