

## Product datasheet for **TA800063BM**

### **clAP2 (BIRC3) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3G4]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI3G4
Applications:	FC, WB
Recommended Dilution:	WB 1:500~2000, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 255-529 of human BIRC3 (NP_001156) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	68.2 kDa
Gene Name:	baculoviral IAP repeat containing 3
Database Link:	<a href="#">NP_001156</a> <a href="#">Entrez Gene 330 Human</a> <a href="#">Q13489</a>



[View online »](#)

**Background:**

This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq, Aug 2011]

**Synonyms:**

AIP1; API2; c-IAP2; CIAP2; HAIP1; HIAP1; MALT2; MIHC; RNF49


**Protein Families:**

Druggable Genome


**Protein Pathways:**

Apoptosis, Focal adhesion, NOD-like receptor signaling pathway, Pathways in cancer, Small cell lung cancer, Ubiquitin mediated proteolysis


**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY BIRC3 (Cat# [RC207764], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BIRC3(Cat# [TA800063]). Positive lysates [LY420093] (100ug) and [LC420093] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-BIRC3 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



HEK293T cells transfected with either [RC207764] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-BIRC3 antibody ([TA800063]), and then analyzed by flow cytometry.