

Product datasheet for TA800063M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

cIAP2 (BIRC3) Mouse Monoclonal Antibody [Clone ID: OTI3G4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3G4
Applications: FC, WB

Recommended Dilution: WB 1:500~2000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 255-529 of human

BIRC3 (NP_001156) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 68.2 kDa

Gene Name: baculoviral IAP repeat containing 3

Database Link: NP 001156

Entrez Gene 330 Human

013489





Background:

This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq, Aug 2011]


Synonyms: AIP1; API2; c-IAP2; CIAP2; HAIP1; HIAP1; MALT2; MIHC; RNF49

Protein Families: Druggable Genome


Protein Pathways: Apoptosis, Focal adhesion, NOD-like receptor signaling pathway, Pathways in cancer, Small

cell lung cancer, Ubiquitin mediated proteolysis

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY BIRC3 (Cat# [RC207764], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BIRC3(Cat# [TA800063]). Positive lysates [LY420093] (100ug) and [LC420093] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-BIRC3 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).





HEK293T cells transfected with either [RC207764] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-BIRC3 antibody ([TA800063]), and then analyzed by flow cytometry.