

Product datasheet for **TA800157AM**

CNOT4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1H2]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1H2
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 190-455 of human CNOT4 (NP_037448) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	70.2 kDa
Gene Name:	CCR4-NOT transcription complex subunit 4
Database Link:	NP_037448 Entrez Gene 53621 Mouse Entrez Gene 312227 Rat Entrez Gene 4850 Human O95628
Background:	The protein encoded by this gene is a subunit of the CCR4-NOT complex, a global transcriptional regulator. The encoded protein interacts with CNOT1 and has E3 ubiquitin ligase activity. Several transcript variants encoding different isoforms have been found for this gene.
Synonyms:	CLONE243; NOT4; NOT4H




[View online »](#)


Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: RNA degradation


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CNOT4 (Cat# [RC217418], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CNOT4(Cat# [TA800157]). Positive lysates [LY415669] (100ug) and [LC415669] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-CNOT4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800157])