

## **Product datasheet for TA800200BM**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SLUG (SNAI2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4E1]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI4E1
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human SNAI2 (NP\_003059) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 29.8 kDa

**Gene Name:** snail family transcriptional repressor 2

Database Link: NP 003059

Entrez Gene 20583 MouseEntrez Gene 25554 RatEntrez Gene 6591 Human

043623

**Background:** This gene encodes a member of the Snail family of C2H2-type zinc finger transcription

factors. The encoded protein acts as a transcriptional repressor that binds to E-box motifs and is also likely to repress E-cadherin transcription in breast carcinoma. This protein is involved in epithelial-mesenchymal transitions and has antiapoptotic activity. Mutations in this gene may be associated with sporatic cases of neural tube defects. [provided by RefSeq]

Synonyms: SLUG; SLUGH1; SNAIL2; WS2D







**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Adherens junction

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SNAI2 ([RC202365], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SNAI2. Positive lysates [LY401070] (100ug) and [LC401070] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC202365] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SNAI2 antibody ([TA800200]), and then analyzed by flow cytometry.