

Product datasheet for TA800325

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RIP (RIPK1) Mouse Monoclonal Antibody [Clone ID: OTI2G3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2G3
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150

Reactivity: Human, Dog

Host: Mouse lgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 133-422 of human

RIPK1(NP_003795) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 75.8 kDa

Gene Name: receptor interacting serine/threonine kinase 1

Database Link: NP 003795

Entrez Gene 102152063 DogEntrez Gene 8737 Human

Q13546

Synonyms: RIP; RIP-1; RIP1

Protein Families: Druggable Genome, Protein Kinase


Protein Pathways: Apoptosis, Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway, Toll-like

receptor signaling pathway







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RIPK1 ([RC216024], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RIPK1. Positive lysates [LY401251] (100ug) and [LC401251] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-RIPK1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-RIPK1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.