

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA800337M

# Factor XIIIa (F13A1) Mouse Monoclonal Antibody [Clone ID: OTI1C9]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI1C9
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human F13A1 (NP_000120) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	79.2 kDa
Gene Name:	coagulation factor XIII A chain
Database Link:	<u>NP_000120</u> <u>Entrez Gene 60327 RatEntrez Gene 74145 MouseEntrez Gene 2162 Human</u> <u>P00488</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### Sactor XIIIa (F13A1) Mouse Monoclonal Antibody [Clone ID: OTI1C9] – TA800337M

Background:This gene encodes the coagulation factor XIII A subunit. Coagulation factor XIII is the last<br/>zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a<br/>heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic<br/>function, and the B subunits do not have enzymatic activity and may serve as plasma carrier<br/>molecules. Platelet factor XIII is comprised only of 2 A subunits, which are identical to those<br/>of plasma origin. Upon cleavage of the activation peptide by thrombin and in the presence of<br/>calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active<br/>enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze<br/>the formation of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus<br/>stabilizing the fibrin clot. It also crosslinks alpha-2-plasmin inhibitor, or fibronectin, to the<br/>alpha chains of fibrin. Factor XIII deficiency is classified into two categories: type I deficiency,<br/>characterized by the lack of both the A and B subunits; and type II deficiency, characterized<br/>by the lack of the A subunit alone. These defects can result in a lifelong bleeding tendency,<br/>defective wound healing, and habitual abortion. [provided by RefSeq, Jul 2008]

Synonyms:	F13A
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Complement and coagulation cascades

### **Product images:**

170	_	
130	-	
100	-	-
70	-	-
55	-	
40	-	
35	-	
25	-	
15	-	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY F13A1 ([RC206464], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-F13A1. Positive lysates [LY400044] (100ug) and [LC400044] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US