

### Product datasheet for TA800365BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ARL2BP Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1G10]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI1G10

Applications: WB

Recommended Dilution: WB: 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARL2BP (NP\_036238) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 18.6 kDa

**Gene Name:** ADP ribosylation factor like GTPase 2 binding protein

Database Link: NP 036238

Entrez Gene 107566 MouseEntrez Gene 498910 RatEntrez Gene 23568 Human

Q9Y2Y0






#### Background:

ADP-ribosylation factor (ARF)-like proteins (ARLs) comprise a functionally distinct group of the ARF family of RAS-related GTPases. The protein encoded by this gene binds to ARL2.GTP with high affinity but does not interact with ARL2.GDP, activated ARF, or RHO proteins. The lack of detectable membrane association of this protein or ARL2 upon activation of ARL2 is suggestive of actions distinct from those of the ARFs. This protein is considered to be the first ARL2-specific effector identified, due to its interaction with ARL2.GTP but lack of ARL2 GTPase-activating protein activity. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.

**Synonyms:** BART; BART1; RP66

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ARL2BP ([RC200198], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARL2BP. Positive lysates [LY415976] (100ug) and [LC415976] (20ug) can be purchased separately from OriGene.