

# **Product datasheet for TA800387M**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SEN1 (MORF4) Mouse Monoclonal Antibody [Clone ID: OTI5E6]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5E6

**Applications:** WB

Recommended Dilution: WB: 1:500
Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-255 of human MORF4

(NP\_006783) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 26.6 kDa

**Gene Name:** mortality factor 4 (pseudogene)

Database Link: NP 006783

Entrez Gene 10934 Human

**Background:** Cellular senescence, the terminal nondividing state that normal cells enter following

completion of their proliferative potential, is the dominant phenotype in hybrids of normal and immortal cells. Fusions of immortal human cell lines with each other have led to their assignment to 1 of several complementation groups. MORF4 is a gene on chromosome 4 that

induces a senescent-like phenotype in cell lines assigned to complementation group B.

[supplied by OMIM]






### SEN1 (MORF4) Mouse Monoclonal Antibody [Clone ID: OTI5E6] - TA800387M

Synonyms: CSR; CSRB; mortality factor 4; SEN; SEN1; senescence (cellular)-related 1; senescence-related,

cellular, 1

**Protein Families:** Transcription Factors

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MORF4 ([RC217344], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MORF4.