

Product datasheet for TA800388M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ARL2BP Mouse Monoclonal Antibody [Clone ID: OTI1F8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1F8
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human, Dog, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARL2BP (NP_036238) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 18.6 kDa

Gene Name: ADP ribosylation factor like GTPase 2 binding protein

Database Link: NP 036238

Entrez Gene 107566 MouseEntrez Gene 498910 RatEntrez Gene 611248 DogEntrez Gene

703999 MonkeyEntrez Gene 23568 Human

O9Y2Y0






Background:


ADP-ribosylation factor (ARF)-like proteins (ARLs) comprise a functionally distinct group of the ARF family of RAS-related GTPases. The protein encoded by this gene binds to ARL2.GTP with high affinity but does not interact with ARL2.GDP, activated ARF, or RHO proteins. The lack of detectable membrane association of this protein or ARL2 upon activation of ARL2 is suggestive of actions distinct from those of the ARFs. This protein is considered to be the first ARL2-specific effector identified, due to its interaction with ARL2.GTP but lack of ARL2 GTPase-activating protein activity. [provided by RefSeq, Jul 2008]

Synonyms: BART; BART1; RP66

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ARL2BP ([RC200198], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARL2BP. Positive lysates [LY415976] (100ug) and [LC415976] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ARL2BP monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).





Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-ARL2BP mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-ARL2BP mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.