

Product datasheet for TA800399M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

A4GNT Mouse Monoclonal Antibody [Clone ID: OTI2H9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2H9
Applications: IHC, WB

Recommended Dilution: WB: 1:500, IHC: 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 121-340 of human

A4GNT (NP_057245) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 39.3 kDa

Gene Name: alpha-1,4-N-acetylglucosaminyltransferase

Database Link: NP 057245

Entrez Gene 51146 Human

Q9UNA3

Background: This gene encodes a protein from the glycosyltransferase 32 family. The enzyme catalyzes the

transfer of N-acetylglucosamine (GlcNAc) to core 2 branched O-glycans. It forms a unique glycan, GlcNAcalpha1-->4Galbeta-->R and is largely associated with the Golgi apparatus

membrane. [provided by RefSeq]


Synonyms: alpha4GnT






Protein Families: Transmembrane

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY A4GNT ([RC221256], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-A4GNT. Positive lysates [LY414153] (100ug) and [LC414153] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-A4GNT mouse monoclonal antibody. ([TA800399]) Dilution: 1:150. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.