

# Product datasheet for TA800407M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Epoxide hydrolase (EPHX1) Mouse Monoclonal Antibody [Clone ID: OTI3F10]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3F10
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 21-230 of human

EPHX1 (NP\_000111) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 52.8 kDa

**Gene Name:** epoxide hydrolase 1

Database Link: NP 000111

Entrez Gene 13849 MouseEntrez Gene 25315 RatEntrez Gene 700180 MonkeyEntrez Gene

<u>2052 Human</u>

P07099





**Background:** Epoxide hydrolase is a critical biotransformation enzyme that converts epoxides from the

degradation of aromatic compounds to trans-dihydrodiols which can be conjugated and excreted from the body. Epoxide hydrolase functions in both the activation and detoxification of epoxides. Mutations in this gene cause preeclampsia, epoxide hydrolase deficiency or increased epoxide hydrolase activity. Alternatively spliced transcript variants encoding the


same protein have been found for this gene. [provided by RefSeq, Dec 2008]

Synonyms: EPHX; EPOX; HYL1; MEH


**Protein Families:** Druggable Genome, Protease

**Protein Pathways:** Metabolism of xenobiotics by cytochrome P450

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY EPHX1 (Cat# [RC200621], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-EPHX1(Cat# [TA800407]). Positive lysates [LY400042] (100ug) and [LC400042] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-EPHX1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).






Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-EPHX1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-EPHX1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-EPHX1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.