

## Product datasheet for **TA800429AM**

### ACADS Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1D2]


#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1D2
Applications:	IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:150
Reactivity:	Human, Monkey, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 34-260 of human ACADS (NP_000008) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	41.7 kDa
Gene Name:	acyl-CoA dehydrogenase, C-2 to C-3 short chain
Database Link:	<a href="#">NP_000008</a> <a href="#">Entrez Gene 11409 Mouse</a> <a href="#">Entrez Gene 64304 Rat</a> <a href="#">Entrez Gene 698196 Monkey</a> <a href="#">Entrez Gene 35 Human</a> <a href="#">P16219</a>
Background:	This gene encodes a tetrameric mitochondrial flavoprotein, which is a member of the acyl-CoA dehydrogenase family. This enzyme catalyzes the initial step of the mitochondrial fatty acid beta-oxidation pathway. Mutations in this gene have been associated with Short Chain Acyl-CoA Dehydrogenase Deficiency. [provided by RefSeq, Jul 2008]




[View online »](#)


**Synonyms:** ACAD3; SCAD  
**Protein Families:** Druggable Genome  
**Protein Pathways:** Butanoate metabolism, Fatty acid metabolism, Metabolic pathways, Valine, leucine and isoleucine degradation

**Product images:**



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-ACADS mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800429])



Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-ACADS mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA800429])



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ACADS (Cat# [RC206855], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACADS (Cat# [TA800429]). Positive lysates [LY400002] (100ug) and [LC400002] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ACADS monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).