

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA800432S

## A4GNT Mouse Monoclonal Antibody [Clone ID: OTI1B10]

#### **Product data:**

| Product Type:           | Primary Antibodies  |
|-------------------------|---|
| Clone Name:             | OTI1B10   |
| Applications:           | WB  |
| Recommended Dilution:   |   |
| Reactivity:             | Human   |
| Host:                   | Mouse   |
| lsotype:                | lgG1  |
| Clonality:              | Monoclonal  |
| Immunogen:              | Human recombinant protein fragment corresponding to amino acids 121-340 of human<br>A4GNT (NP_057245) produced in E.coli.   |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.  |
| Concentration:          | 1 mg/ml   |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography<br>(protein A/G)  |
| Conjugation:            | Unconjugated  |
| Storage:                | Store at -20°C as received.   |
| Stability:              | Stable for 12 months from date of receipt.  |
| Predicted Protein Size: | 39.3 kDa  |
| Gene Name:              | alpha-1,4-N-acetylglucosaminyltransferase   |
| Database Link:          | <u>NP_057245</u><br><u>Entrez Gene 51146 Human</u><br><u>Q9UNA3</u>   |
| Background:             | This gene encodes a protein from the glycosyltransferase 32 family. The enzyme catalyzes the transfer of N-acetylglucosamine (GlcNAc) to core 2 branched O-glycans. It forms a unique glycan, GlcNAcalpha1>4Galbeta>R and is largely associated with the Golgi apparatus membrane. [provided by RefSeq] |
| Synonyms:               | alpha4GnT   |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Protein Families: Trai

Transmembrane

## **Product images:**

| 170 | -  |   |
|-----|----|---|
| 130 | -  |   |
| 100 | -  |   |
| 70  | -  |   |
| 55  | -  |   |
| 40  | -  | - |
| 35  | -  |   |
| 25  | -  |   |
| 15  | -  |   |
| 10  | -1 |   |
|     |    |   |

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY A4GNT ([RC221256], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-A4GNT. Positive lysates [LY414153] (100ug) and [LC414153] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US