

## Product datasheet for TA800501BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### PPT1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1B10]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1B10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Dog, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 100-306 of human PPT1

(NP\_000301) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 31.2 kDa

**Gene Name:** palmitoyl-protein thioesterase 1

Database Link: NP 000301

Entrez Gene 475316 DogEntrez Gene 693377 MonkeyEntrez Gene 5538 Human

P50897

**Background:** The protein encoded by this gene is a small glycoprotein involved in the catabolism of lipid-

modified proteins during lysosomal degradation. The encoded enzyme removes thioester-linked fatty acyl groups such as palmitate from cysteine residues. Defects in this gene are a cause of infantile neuronal ceroid lipofuscinosis 1 (CLN1, or INCL) and neuronal ceroid lipofuscinosis 4 (CLN4). Two transcript variants encoding different isoforms have been found

for this gene. [provided by RefSeq, Dec 2008]






Synonyms: CLN1; INCL; PPT


**Protein Families:** Druggable Genome

**Protein Pathways:** Fatty acid elongation in mitochondria, Lysosome, Metabolic pathways

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PPT1 ([RC203278], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PPT1. Positive lysates [LY400121] (100ug) and [LC400121] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PPT1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).