

# **Product datasheet for TA800573AM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### C2ORF25 (MMADHC) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1G4]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1G4

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 39-296 of human

MMADHC (NP\_056517) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 32.8 kDa

**Gene Name:** methylmalonic aciduria and homocystinuria, cblD type

Database Link: NP 056517

Entrez Gene 109129 MouseEntrez Gene 362134 RatEntrez Gene 27249 Human

O9H3L0

**Background:** This gene encodes a mitochondrial protein that is involved in an early step of vitamin B12

metabolism. Vitamin B12 (cobalamin) is essential for normal development and survival in humans. Mutations in this gene cause methylmalonic aciduria and homocystinuria type cblD (MMADHC), a disorder of cobalamin metabolism that is characterized by decreased levels of the coenzymes adenosylcobalamin and methylcobalamin. Pseudogenes have been identified


on chromosomes 11 and X. [provided by RefSeq, Nov






Synonyms: C2orf25; cblD; CL25022

## **Product images:**





HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY MMADHC (Cat# [RC204801], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MMADHC(Cat# [TA800573]). Positive lysates [LY414395] (100ug) and [LC414395] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-MMADHC monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).