

## **Product datasheet for TA800679**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TRIF (TICAM1) Mouse Monoclonal Antibody [Clone ID: OTI1G7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1G7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 120-420 of human

TICAM1 (NP\_891549) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 76.2 kDa

**Gene Name:** toll like receptor adaptor molecule 1

Database Link: NP 891549

Entrez Gene 574294 MonkeyEntrez Gene 148022 Human

O8IUC6





#### Background:


This gene encodes an adaptor protein containing a Toll/interleukin-1 receptor (TIR) homology domain, which is an intracellular signaling domain that mediates protein-protein interactions between the Toll-like receptors (TLRs) and signal-transduction components. This protein is involved in native immunity against invading pathogens. It specifically interacts with toll-like receptor 3, but not with other TLRs, and this association mediates dsRNA induction of interferon-beta through activation of nuclear factor kappa-B, during an antiviral immune response. [provided by RefSeq, Jan 2012]


Synonyms: IIAE6; MyD88-3; PRVTIRB; TICAM-1; TRIF

**Protein Families:** Druggable Genome

**Protein Pathways:** Toll-like receptor signaling pathway

# **Product images:**





HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TICAM1 (Cat# [RC217522], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TICAM1(Cat# TA800679). Positive lysates [LY403651] (100ug) and [LC403651] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-TICAM1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).