

# Product datasheet for TA800701BM

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ALK Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1F1]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1F1

**Applications:** LMNX, WB **Recommended Dilution:** WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1060-1620 of human

ALK (NP\_004295) produced in SF9 cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 176.3 kDa

**Gene Name:** anaplastic lymphoma receptor tyrosine kinase

Database Link: NP 004295

Entrez Gene 11682 MouseEntrez Gene 266802 RatEntrez Gene 238 Human

O9UM73






#### Background:

This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor superfamily. This protein comprises an extracellular domain, an hydrophobic stretch corresponding to a single pass transmembrane region, and an intracellular kinase domain. It plays an important role in the development of the brain and exerts its effects on specific neurons in the nervous system. This gene has been found to be rearranged, mutated, or amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma, and non-small cell lung cancer. The chromosomal rearrangements are the most common genetic alterations in this gene, which result in creation of multiple fusion genes in tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 2), ALK/RANBP2 (chromosome 2), ALK/ATIC (chromosome 2), ALK/TFG (chromosome 3), ALK/NPM1 (chromosome 5), ALK/SQSTM1 (chromosome 5), ALK/KIF5B (chromosome 10), ALK/CLTC (chromosome 17), ALK/TPM4 (chromosome 19), and ALK/MSN (chromosome X). [provided by RefSeq, Jan 2011]

Synonyms: CD246; NBLST3


**Protein Families:** Druggable Genome, Protein Kinase

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALK ([RC222485], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALK. Positive lysates [LY418072] (100ug) and [LC418072] (20ug) can be purchased separately from OriGene.





ALK Luminex with 1D7 Capture ([TA800796]) and 1F1 Detection ([TA800701]) Antibodies. Substrate used: full length HEK293 cells expressed recombinant ALK protein ([TP322485]).