

A TOOR RESEARCH

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA800795M

ALK Mouse Monoclonal Antibody [Clone ID: OTI5C11]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5C11
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ALK (NP_004295) produced in HEK293T cell
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	176.3 kDa
Gene Name:	anaplastic lymphoma receptor tyrosine kinase
Database Link:	<u>NP_004295</u> <u>Entrez Gene 11682 MouseEntrez Gene 266802 RatEntrez Gene 238 Human</u> <u>Q9UM73</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Scheme Constant ALK Mouse Monoclonal Antibody [Clone ID: OTI5C11] – TA800795M


Background:This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor
superfamily. This protein comprises an extracellular domain, an hydrophobic stretch
corresponding to a single pass transmembrane region, and an intracellular kinase domain. It
plays an important role in the development of the brain and exerts its effects on specific
neurons in the nervous system. This gene has been found to be rearranged, mutated, or
amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma,
and non-small cell lung cancer. The chromosomal rearrangements are the most common
genetic alterations in this gene, which result in creation of multiple fusion genes in
tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 3), ALK/RANBP2
(chromosome 2), ALK/ATIC (chromosome 2), ALK/KIF5B (chromosome 10), ALK/CLTC
(chromosome 17), ALK/TPM4 (chromosome 19), and ALK/MSN (chromosome X). [provided by
RefSeq, Jan 2011]

Synonyms: CD246; NBLST3

Protein Families:

Druggable Genome, Protein Kinase

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALK ([RC222485], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALK. Positive lysates [LY418072] (100ug) and [LC418072] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US