

Product datasheet for TA800805

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CTNND1 Mouse Monoclonal Antibody [Clone ID: OTI2G7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2G7
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CTNND1 (NP_001078927) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 108 kDa

Gene Name: catenin delta 1

Database Link: NP 001078927

Entrez Gene 12388 MouseEntrez Gene 311163 RatEntrez Gene 1500 Human

060716

Background: This gene encodes a member of the Armadillo protein family, which function in adhesion

between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane

protein 2 (TMX2) gene. [provided by RefSeq, Dec 2010]




Synonyms: CAS; CTNND; p120; p120(CAS); p120(CTN); P120CAS; P120CTN


Protein Families: Druggable Genome

Protein Pathways: Adherens junction, Leukocyte transendothelial migration

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CTNND1 ([RC222771], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CTNND1. Positive lysates [LY421307] (100ug) and [LC421307] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from different cell line by using anti-CTNND1 monoclonal antibody (1:500).







Equivalent amounts of cell lysates (10 ug per lane) of wild-type Hela cells (WT, Cat# LC810HELA) and CTNND1-Knockout Hela cells (KO, Cat# [LC810057]) were separated by SDS-PAGE and immunoblotted with anti-CTNND1 monoclonal antibody TA800805. Then the blotted membrane was stripped and reprobed with anti-b-actin antibody ([TA811000]) as a loading control (1:500).

Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-CTNND1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800805) (1:150)