

Product datasheet for **TA800824BM**

CD99 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1D5]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1D5
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CD99 (NP_002405) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	16.7 kDa
Gene Name:	CD99 molecule
Database Link:	NP_002405 Entrez Gene 4267 Human P14209



[View online »](#)

Background:

The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. Cyclophilin A binds to CD99 and may act as a signaling regulator of CD99. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Synonyms:


HBA71; MIC2; MIC2X; MIC2Y; MSK5X

Protein Families:


Druggable Genome, Transmembrane

Protein Pathways:


Cell adhesion molecules (CAMs), Leukocyte transendothelial migration

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CD99 (Cat# [RC204056], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD99 (Cat# [TA800824]). Positive lysates [LY400863] (100ug) and [LC400863] (20ug) can be purchased separately from OriGene.




Western blot analysis of A549 cell lysate (35ug) by using anti-CD99 monoclonal antibody. Dilution: 1:500




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-CD99 mouse monoclonal antibody. ([TA800824]) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-CD99 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA800824])



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-CD99 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA800824])



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-CD99 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA800824])



Flow cytometric Analysis of living Jurkat cells, using anti-CD99 antibody ([TA800824]), (Red), compared to a nonspecific negative control antibody, (Blue).