

### Product datasheet for TA800856AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# MGMT Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI4E6]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4E6
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human MGMT (NP\_002403) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: O-6-methylguanine-DNA methyltransferase

Database Link: NP 002403

Entrez Gene 4255 Human

P16455

Synonyms: methylguanine-DNA methyltransferase; O-6-methylguanine-DNA methyltransferase; O6-


methylguanine-DNA methyltransferase; OTTHUMP0000020741

**Protein Families:** Druggable Genome







# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MGMT ([RC229131], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MGMT. Positive lysates [LY432155] (100ug) and [LC432155] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-MGMT monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human). Dilution: 1:2000




Western blot analysis of extracts (35ug) from 5 differentcell lines by using anti-MGMT monoclonal antibody (1:500).






Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-MGMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA800856])



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-MGMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA800856])



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-MGMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA800856])