

Product datasheet for TA800883M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PCNA Mouse Monoclonal Antibody [Clone ID: OTI5F9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5F9
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PCNA (NP_002583) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 28.6 kDa

Gene Name: proliferating cell nuclear antigen

Database Link: NP 002583

Entrez Gene 18538 MouseEntrez Gene 25737 RatEntrez Gene 5111 Human

P12004

Background: The protein encoded by this gene is found in the nucleus and is a cofactor of DNA

polymerase delta. The encoded protein acts as a homotrimer and helps increase the

processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome. [provided by

RefSeq, Jul 2008]






Synonyms: ATLD2


Protein Families: Druggable Genome, Stem cell - Pluripotency

Protein Pathways: Base excision repair, Cell cycle, DNA replication, Mismatch repair, Nucleotide excision repair

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PCNA ([RC201741], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PCNA. Positive lysates [LY400929] (100ug) and [LC400929] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PCNA monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).