

Product datasheet for TA800923M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD99 Mouse Monoclonal Antibody [Clone ID: OTI1A1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A1
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD99 (NP_002405) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 18.85 kDa

Gene Name: CD99 molecule

Database Link: NP 002405

Entrez Gene 4267 Human

P14209


Synonyms: HBA71; MIC2; MIC2X; MIC2Y; MSK5X

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: Cell adhesion molecules (CAMs), Leukocyte transendothelial migration



Product images:




Western blot analysis of extracts (35ug) from Jurkat and HUVEC cell lines by using anti-KRT14 monoclonal antibody (1:500).




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-CD99 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-CD99 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min



Immunohistochemical staining of paraffinembedded Human appendix tissue within the normal limits using anti-CD99 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.