

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA801186S

# p18 INK4c (CDKN2C) Mouse Monoclonal Antibody [Clone ID: OTI1E4]

## **Product data:**

Product Type:	Primary Antibodies	
Clone Name:	OTI1E4	
Applications:	WB	
Recommended Dilution:	WB 1:2000	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human CDKN2C (NP_001253) produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	17.9 kDa	
Gene Name:	cyclin-dependent kinase inhibitor 2C	
Database Link:	<u>NP_001253</u> Entrez Gene 12580 MouseEntrez Gene 54238 RatEntrez Gene 1031 Human P42773	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	p18 INK4c (CDKN2C) Mouse Monoclonal Antibody [Clone ID: OTI1E4] – TA801186S			
Background:	The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase inhibitors. This protein has been shown to interact with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. Ectopic expression of this gene was shown to suppress the growth of human cells in a manner that appears to correlate with the presence of a wild-type RB1 function. Studies in the knockout mice suggested the roles of this gene in regulating spermatogenesis, as well as in suppressing tumorigenesis. Two alternatively spliced transcript variants of this gene, which encode an identical protein, have been reported. [provided by RefSeq, Jul 2008]			
Synonyms:	INK4C; p18; p18-INK4C			
Protein Families:	Druggable Genome			
Protein Pathway	s: Cell cycle			

## **Product images:**

170	-	
130	-	
100	_	
70	_	
55	_	
40	-	
35	-	
25	-	
15	-	
10	_	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CDKN2C ([RC213083], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CDKN2C. Positive lysates [LY420041] (100ug) and [LC420041] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US