

# Product datasheet for TA801262S

## ALK Mouse Monoclonal Antibody [Clone ID: OTI2H8]

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Clone Name:	OTI2H8
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1300-1620 of human ALK (NP_004295) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	176.3 kDa
Gene Name:	anaplastic lymphoma receptor tyrosine kinase
Database Link:	<u>NP_004295</u> <u>Entrez Gene 11682 MouseEntrez Gene 266802 RatEntrez Gene 238 Human</u> <u>Q9UM73</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### Serigene ALK Mouse Monoclonal Antibody [Clone ID: OTI2H8] – TA801262S

Background:This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor<br/>superfamily. This protein comprises an extracellular domain, an hydrophobic stretch<br/>corresponding to a single pass transmembrane region, and an intracellular kinase domain. It<br/>plays an important role in the development of the brain and exerts its effects on specific<br/>neurons in the nervous system. This gene has been found to be rearranged, mutated, or<br/>amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma,<br/>and non-small cell lung cancer. The chromosomal rearrangements are the most common<br/>genetic alterations in this gene, which result in creation of multiple fusion genes in<br/>tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 3), ALK/RANBP2<br/>(chromosome 2), ALK/ATIC (chromosome 2), ALK/KIF5B (chromosome 10), ALK/CLTC<br/>(chromosome 17), ALK/TPM4 (chromosome 19), and ALK/MSN (chromosome X). [provided by<br/>RefSeq, Jan 2011]

Synonyms: CD246; NBLST3

Protein Families: Druggable Genome, Protein Kinase

#### **Product images:**

170	-	
130		
100		-
70		
55		
40	-	
35		
25	-	
15	-1	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALK ([RC222485], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALK. Positive lysates [LY418072] (100ug) and [LC418072] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US