

Product datasheet for **TA801287AM**

ALK Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1A4
Applications:	IHC, LMNX, WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1300-1620 of human ALK (NP_004295) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	176.3 kDa
Gene Name:	anaplastic lymphoma receptor tyrosine kinase
Database Link:	NP_004295 Entrez Gene 238 Human Q9UM73



[View online »](#)

Background:


This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor superfamily. This protein comprises an extracellular domain, an hydrophobic stretch corresponding to a single pass transmembrane region, and an intracellular kinase domain. It plays an important role in the development of the brain and exerts its effects on specific neurons in the nervous system. This gene has been found to be rearranged, mutated, or amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma, and non-small cell lung cancer. The chromosomal rearrangements are the most common genetic alterations in this gene, which result in creation of multiple fusion genes in tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 2), ALK/RANBP2 (chromosome 2), ALK/ATIC (chromosome 2), ALK/TFG (chromosome 3), ALK/NPM1 (chromosome 5), ALK/SQSTM1 (chromosome 5), ALK/KIF5B (chromosome 10), ALK/CLTC (chromosome 17), ALK/TPM4 (chromosome 19), and ALK/MSN (chromosome X). [provided by RefSeq, Jan 2011]

Synonyms:


CD246; NBLST3

Protein Families:


Druggable Genome, Protein Kinase

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALK ([RC222485], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALK. Positive lysates [LY418072] (100ug) and [LC418072] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from H2228 and A549 cell lines by using anti-ALK monoclonal antibody. ([TA801287], 1:10,000)




Immunohistochemical staining of paraffin-embedded Human non-small cell lung cancer sample with ALK translocation detected by FISH using anti-ALK mouse monoclonal antibody. ([TA801287], 1:50; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Human large B cell lymphoma with ALK translocation using anti-ALK mouse monoclonal antibody. ([TA801287], 1:50; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded ALK-positive lung tumor xenograft using anti-ALK mouse monoclonal antibody. ([TA801287], 1:50; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.0, 120°C for 3min)



Immunohistochemical staining of paraffin-embedded Human non-small cell lung cancer sample with EML4-ALK translocation detected by PCR using anti-ALK mouse monoclonal antibody. ([TA801287], 1:50; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.0, 120°C for 3min)



Immunohistochemistry staining of paraffin-embedded human cell line H1975 (upper left) and H2228 (upper right) on IHC antibody quality control slide using anti-ALK mouse monoclonal antibody [TA801287] (1:400). The ALK rearrangement in H2228 cells is labeled with ALK Breakapart probe in FISH test (lower right, 60X) and the control of H1975 cell at the same FISH probe test (lower left, 60X).



Immunohistochemical staining of paraffin-embedded human ALK-positive lung cancer tissue using anti-ALK mouse monoclonal antibody. ([TA801287], 1:100 for 30 min at RT; heat-induced epitope retrieval by TEE, pH9.0)






Figure from citation: Immunohistochemistry of ALK protein level by using anti-ALK antibody in human right cervical lymph node. Dilution: 1:200
[View Citation](#)



ALK Luminex with 15B2 Capture ([TA801288]) and 1A4 Detection ([TA801287]) Antibodies. Substrate used: full length HEK293 cells expressed recombinant ALK protein ([TP322485]).



ALK Luminex with 10D7 Capture ([TA801306]) and 1A4 Detection ([TA801287]) Antibodies. Substrate used: full length HEK293 cells expressed recombinant ALK protein ([TP322485]).