

Product datasheet for TA801482BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PI 3 Kinase catalytic subunit alpha (PIK3CA) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI7A7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7A7

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 303-631 of human

PIK3CA (NP 006209) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 124.1 kDa

Gene Name: phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha

Database Link: NP 006209

Entrez Gene 18706 MouseEntrez Gene 170911 RatEntrez Gene 5290 Human

P42336

Background: Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa

catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. [provided by RefSeq, Jul 2008]





PI 3 Kinase catalytic subunit alpha (PIK3CA) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI7A7] – TA801482BM

Synonyms: CLOVE; CWS5; MCAP; MCM; MCMTC; p110-alpha; PI3K; PI3K-alpha

Protein Families: Druggable Genome

Protein Pathways: Acute myeloid leukemia, Apoptosis, B cell receptor signaling pathway, Chemokine signaling


pathway, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Glioma, Inositol phosphate metabolism, Insulin signaling pathway, Jak-STAT signaling pathway, Leukocyte transendothelial migration, Melanoma, mTOR signaling

pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Phosphatidylinositol signaling system, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of actin cytoskeleton,

Renal cell carcinoma, Small cell lung cancer, T cell receptor signaling pathway, Toll-like

receptor signaling pathway, Type II diabetes mellitus, VEGF signaling pathway

Product images:



Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-PIK3CA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA801482])