

### Product datasheet for TA801631M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Dicer (DICER1) Mouse Monoclonal Antibody [Clone ID: OTI1E11]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1E11

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1666-1922 of human

DICER1 (NP\_085124) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 218.5 kDa

**Gene Name:** dicer 1, ribonuclease III

Database Link: NP 085124

Entrez Gene 192119 MouseEntrez Gene 299284 RatEntrez Gene 23405 Human

O9UPY3

Background: This gene encodes a protein possessing an RNA helicase motif containing a DEXH box in its


amino terminus and an RNA motif in the carboxy terminus. The encoded protein functions as a ribonuclease and is required by the RNA interference and small temporal RNA (stRNA) pathways to produce the active small RNA component that represses gene expression. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep




Synonyms: DCR1; Dicer; Dicer1e; HERNA; K12H4.8-LIKE; MNG1; RMSE2

**Protein Families:** Druggable Genome

# **Product images:**



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-DICER1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-DICER1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.