

## Product datasheet for **TA801644M**

### Dicer (DICER1) Mouse Monoclonal Antibody [Clone ID: OTI1G8]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI1G8
Applications:	IHC, WB
Recommended Dilution:	IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1666-1922 of human DICER1 (NP_085124) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	218.5 kDa
Gene Name:	dicer 1, ribonuclease III
Database Link:	<a href="#">NP_085124</a> <a href="#">Entrez Gene 192119 Mouse</a> <a href="#">Entrez Gene 299284 Rat</a> <a href="#">Entrez Gene 23405 Human</a> <a href="#">Q9UPY3</a>
Background:	This gene encodes a protein possessing an RNA helicase motif containing a DEXH box in its amino terminus and an RNA motif in the carboxy terminus. The encoded protein functions as a ribonuclease and is required by the RNA interference and small temporal RNA (stRNA) pathways to produce the active small RNA component that represses gene expression. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep


[View online »](#)


**Synonyms:** DCR1; Dicer; Dicer1e; HERNA; K12H4.8-LIKE; MNG1; RMSE2

**Protein Families:** Druggable Genome


**Product images:**



Western blot analysis of HT29 cell lysate (35ug)  
by using anti-DICER1 monoclonal antibody.  
Dilution: 1:500



Immunohistochemical staining of paraffin-  
embedded Adenocarcinoma of Human ovary  
tissue using anti-DICER1 mouse monoclonal  
antibody. Heat-induced epitope retrieval by EDTA  
solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-DICER1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.