

Product datasheet for TA801668M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Neuraminidase (NEU1) Mouse Monoclonal Antibody [Clone ID: OTI3C4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C4

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 48-315 of human NEU1

(NP_000425)produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 40.2 kDa

Gene Name: neuraminidase 1 (lysosomal sialidase)

Database Link: NP 000425

Entrez Gene 4758 Human

099519



Neuraminidase (NEU1) Mouse Monoclonal Antibody [Clone ID: OTI3C4] - TA801668M

Background: The protein encoded by this gene is a lysosomal enzyme that cleaves terminal sialic acid

residues from substrates such as glycoproteins and glycolipids. In the lysosome, this enzyme is part of a heterotrimeric complex together with beta-galactosidase and cathepsin A (the latter is also referred to as 'protective protein'). Mutations in this gene can lead to sialidosis, a lysosomal storage disease that can be type 1 (cherry red spot-myoclonus syndrome or normosomatic type), which is late-onset, or type 2 (the dysmorphic type), which occurs at an


earlier age with increased severity. [provided by RefSeq, Jul 2008]

Synonyms: NANH; NEU; SIAL1

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Lysosome, Other glycan degradation, Sphingolipid metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NEU1 ([RC200386], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEU1. Positive lysates [LY424720] (100ug) and [LC424720] (20ug) can be purchased separately from OriGene.