

## Product datasheet for TA801670M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **Neuraminidase (NEU1) Mouse Monoclonal Antibody [Clone ID: OTI3F5]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3F5

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 48-315 of human NEU1

(NP\_000425)produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 40.2 kDa

**Gene Name:** neuraminidase 1 (lysosomal sialidase)

Database Link: NP 000425

Entrez Gene 4758 Human

099519



#### Neuraminidase (NEU1) Mouse Monoclonal Antibody [Clone ID: OTI3F5] - TA801670M

**Background:** The protein encoded by this gene is a lysosomal enzyme that cleaves terminal sialic acid

residues from substrates such as glycoproteins and glycolipids. In the lysosome, this enzyme is part of a heterotrimeric complex together with beta-galactosidase and cathepsin A (the latter is also referred to as 'protective protein'). Mutations in this gene can lead to sialidosis, a lysosomal storage disease that can be type 1 (cherry red spot-myoclonus syndrome or normosomatic type), which is late-onset, or type 2 (the dysmorphic type), which occurs at an


earlier age with increased severity. [provided by RefSeq, Jul 2008]

Synonyms: NANH; NEU; SIAL1

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Lysosome, Other glycan degradation, Sphingolipid metabolism

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NEU1 ([RC200386], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEU1. Positive lysates [LY424720] (100ug) and [LC424720] (20ug) can be purchased separately from OriGene.