

Product datasheet for TA801709AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PI 3 Kinase Class 2A (PIK3C2A) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2F10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 230-560 of human

PIK3C2A (NP 002636) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 190.5 kDa

Gene Name: phosphatidylinositol-4-phosphate 3-kinase catalytic subunit type 2 alpha

Database Link: NP 002636

Entrez Gene 18704 MouseEntrez Gene 5286 Human

000443





PI 3 Kinase Class 2A (PIK3C2A) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2F10] – TA801709AM

Background: The protein encoded by this gene belongs to the phosphoinositide 3-kinase (PI3K) family. PI3-

kinases play roles in signaling pathways involved in cell proliferation, oncogenic

transformation, cell survival, cell migration, and intracellular protein trafficking. This protein contains a lipid kinase catalytic domain as well as a C-terminal C2 domain, a characteristic of class II PI3-kinases. C2 domains act as calcium-dependent phospholipid binding motifs that mediate translocation of proteins to membranes, and may also mediate protein-protein interactions. The PI3-kinase activity of this protein is not sensitive to nanomolar levels of the inhibitor wortmanin. This protein was shown to be able to be activated by insulin and may be


involved in integrin-dependent signaling. [provided by RefSeq, Jul 2008]

Synonyms: CPK; PI3-K-C2(ALPHA); PI3-K-C2A

Protein Families: Druggable Genome

Protein Pathways: Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system

Product images:



E.coli lysate (left lane) and E.coli lysate expressing human recombinant protein fragment corresponding to amino acids 230-560 of human PIK3C2A (NP_002636) were separated by SDS-PAGE and immunoblotted with anti-PIK3C2A.