

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA801709M

# PI 3 Kinase Class 2A (PIK3C2A) Mouse Monoclonal Antibody [Clone ID: OTI2F10]

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI2F10
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 230-560 of human PIK3C2A (NP_002636) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	190.5 kDa
Gene Name:	phosphatidylinositol-4-phosphate 3-kinase catalytic subunit type 2 alpha
Database Link:	<u>NP_002636</u> <u>Entrez Gene 18704 MouseEntrez Gene 5286 Human</u> <u>O00443</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	3 Kinase Class 2A (PIK3C2A) Mouse Monoclonal Antibody [Clone ID: OTI2F10] – TA801709M		
Background:	The protein encoded by this gene belongs to the phosphoinositide 3-kinase (PI3K) family. PI3 kinases play roles in signaling pathways involved in cell proliferation, oncogenic transformation, cell survival, cell migration, and intracellular protein trafficking. This protein contains a lipid kinase catalytic domain as well as a C-terminal C2 domain, a characteristic of class II PI3-kinases. C2 domains act as calcium-dependent phospholipid binding motifs that mediate translocation of proteins to membranes, and may also mediate protein-protein interactions. The PI3-kinase activity of this protein is not sensitive to nanomolar levels of the inhibitor wortmanin. This protein was shown to be able to be activated by insulin and may be involved in integrin-dependent signaling. [provided by RefSeq, Jul 2008]		
Synonyms:	СРК; РІЗ-К-С2(АLPHA); РІЗ-К-С2А		
<b>Protein Families</b>	: Druggable Genome		
Protein Pathway	<i>is:</i> Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system		

## **Product images:**

170	_	
130	_	
100	_	
70	_	
55	_	
40	_	
35	_	
25	-	
15	_	
10	-	

E.coli lysate (left lane) and E.coli lysate expressing human recombinant protein fragment corresponding to amino acids 230-560 of human PIK3C2A (NP\_002636) were separated by SDS-PAGE and immunoblotted with anti-PIK3C2A.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US