

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA801727AM

Neuraminidase (NEU1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3D4]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI3D4	
Applications:	IHC, WB	
Recommended Dilution:	WB 1:2000, IHC 1:150	
Reactivity:	Human	
Host:	Mouse	
lsotype:	lgG2a	
Clonality:	Monoclonal	
Immunogen:	Human recombinant protein fragment corresponding to amino acids 48-315 of human NEU1 (NP_000425)produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	0.5 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Biotin	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	40.2 kDa	
Gene Name:	neuraminidase 1 (lysosomal sialidase)	
Database Link:	<u>NP_000425</u> <u>Entrez Gene 4758 Human</u> <u>Q99519</u>	




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Neuraminidase (NEU1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3D4] – TA801727AM	
Background:	The protein encoded by this gene is a lysosomal enzyme that cleaves terminal sialic acid residues from substrates such as glycoproteins and glycolipids. In the lysosome, this enzyme is part of a heterotrimeric complex together with beta-galactosidase and cathepsin A (the latter is also referred to as 'protective protein'). Mutations in this gene can lead to sialidosis, a lysosomal storage disease that can be type 1 (cherry red spot-myoclonus syndrome or normosomatic type), which is late-onset, or type 2 (the dysmorphic type), which occurs at an earlier age with increased severity. [provided by RefSeq, Jul 2008]	
Synonyms:	NANH; NEU; SIAL1	
Protein Families	: Druggable Genome, Transmembrane	
Protein Pathway	<i>ys:</i> Lysosome, Other glycan degradation, Sphingolipid metabolism	

Product images:

170	-	
130		
100		
70		
55		
40	-	
35		
25	-	
15	-1	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NEU1 (Cat# [RC200386], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEU1(Cat# [TA801727]). Positive lysates [LY424720] (100ug) and [LC424720] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Neuraminidase (NEU1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3D4] – TA801727AM




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US






Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-NEU1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA801727])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US