

Product datasheet for TA801748S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD137 (TNFRSF9) Mouse Monoclonal Antibody [Clone ID: OTI5H5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5H5

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human

Host: Mouse

Isotype: IgG1
Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 24-186 of human

TNFRSF9 (NP 001552) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: tumor necrosis factor receptor superfamily member 9

Database Link: NP 001552

Entrez Gene 3604 Human

Q07011

Background: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor

contributes to the clonal expansion, survival, and development of T cells. It can also induce

proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 co-stimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-

kappaB. [provided by RefSeq, Jul 2008]




CD137 (TNFRSF9) Mouse Monoclonal Antibody [Clone ID: OTI5H5] - TA801748S

Synonyms: 4-1BB; CD137; CDw137; ILA

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Cytokine-cytokine receptor interaction

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TNFRSF9 ([RC200664], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNFRSF9.