

Product datasheet for TA801830S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD23 (FCER2) Mouse Monoclonal Antibody [Clone ID: OTI9B6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI9B6
Applications: FC, WB

Reactivity: WB 1:2000 Human

Host: Mouse IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 48-321 of human FCER2

(NP_001993) produced in SF9 cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 36.3 kDa

Gene Name: Fc fragment of IgE receptor II

Database Link: NP 001993

Entrez Gene 2208 Human

P06734

Background: The protein encoded by this gene is a B-cell specific antigen, and a low-affinity receptor for

IgE. It has essential roles in B cell growth and differentiation, and the regulation of IgE production. This protein also exists as a soluble secreted form, then functioning as a potent mitogenic growth factor. Alternatively spliced transcript variants encoding different isoforms

have been described for this gene. [provided by RefSeq, Jul




Synonyms: BLAST-2; CD23; CD23A; CLEC4J; FCE2; IGEBF


Protein Families: Secreted Protein, Transmembrane

Protein Pathways: Hematopoietic cell lineage

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FCER2 ([RC204335], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FCER2 (1:2000).




Western blot analysis of extracts (35ug) from 2 cell lines lysates by using anti-FCER2 monoclonal antibody (1:500).









Flow cytometric analysis of living RPMI-8226 cells, using anti-FCER2 antibody ([TA801830], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).