

## **Product datasheet for TA802092BM**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## VILIP1 (VSNL1) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI7E12]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI7E12
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 2-191 of human VSNL1

(NP\_003376) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 22 kDa

**Gene Name:** visinin like 1 **Database Link:** NP 003376

Entrez Gene 24877 RatEntrez Gene 26950 MouseEntrez Gene 7447 Human

P62760

**Background:** This gene is a member of the visinin/recoverin subfamily of neuronal calcium sensor

proteins. The encoded protein is strongly expressed in granule cells of the cerebellum where it associates with membranes in a calcium-dependent manner and modulates intracellular signaling pathways of the central nervous system by directly or indirectly regulating the activity of adenylyl cyclase. Alternatively spliced transcript variants have been observed, but

their full-length nature has not been determined. [provided by RefSeq, Jul 2008]







Synonyms: HLP3; HPCAL3; HUVISL1; VILIP; VILIP-1

**Protein Families:** Druggable Genome


## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY VSNL1 ([RC205337], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-VSNL1. Positive lysates [LY418720] (100ug) and [LC418720] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human adult heart tissue using anti-VSNL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802092])



Immunohistochemical staining of paraffinembedded Human embryonic brain cortex tissue using anti-VSNL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802092])





Immunohistochemical staining of paraffinembedded Human embryonic cerebellum using anti-VSNL1 mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802092])