

Product datasheet for TA802150S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Mitofusin 1 (MFN1) Mouse Monoclonal Antibody [Clone ID: OTI4H4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4H4
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 209-469 of human

MFN1 (NP_284941) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 84 kDa

Gene Name: mitofusin 1

Database Link: NP 284941

Entrez Gene 67414 MouseEntrez Gene 192647 RatEntrez Gene 55669 Human

08IWA4

Background: The protein encoded by this gene is a mediator of mitochondrial fusion. This protein and

mitofusin 2 are homologs of the Drosophila protein fuzzy onion (Fzo). They are mitochondrial

membrane proteins that interact with each other to facilitate mitochondrial targeting.


[provided by RefSeq, Jul 2008]

Synonyms: hfzo1; hfzo2







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MFN1 ([RC207184], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MFN1. Positive lysates [LY403252] (100ug) and [LC403252] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-MFN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802150])



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-MFN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802150])