

Product datasheet for TA802187M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OXSM Mouse Monoclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 78-343 of human OXSM

(NP_060367) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 48.7 kDa

Database Link: NP 060367

Entrez Gene 71147 MouseEntrez Gene 289934 Rat

Background: This gene encodes a beta-ketoacyl synthetase. The encoded enzyme is required for

elongation of fatty acid chains in the mitochondria. Alternatively spliced transcript variants

have been described. [provided by RefSeq, Feb 2009]


Synonyms: FASN2D; KASI; KS
Protein Families: Druggable Genome

Protein Pathways: Fatty acid biosynthesis, Metabolic pathways






Product images:




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-OXSM mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-OXSM mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-OXSM mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.