

# Product datasheet for TA802207M

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## DDIT3 Mouse Monoclonal Antibody [Clone ID: OTI4C4]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI4C4

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human DDIT3 (NP\_004074) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 19 kDa

**Gene Name:** DNA damage inducible transcript 3

Database Link: NP 004074

Entrez Gene 13198 MouseEntrez Gene 29467 RatEntrez Gene 1649 Human

P35638





#### Background:


This gene encodes a member of the CCAAT/enhancer-binding protein (C/EBP) family of transcription factors. The protein functions as a dominant-negative inhibitor by forming heterodimers with other C/EBP members, such as C/EBP and LAP (liver activator protein), and preventing their DNA binding activity. The protein is implicated in adipogenesis and erythropoiesis, is activated by endoplasmic reticulum stress, and promotes apoptosis. Fusion of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by translocation generates chimeric proteins in myxoid liposarcomas or Ewing sarcoma. Multiple alternatively spliced transcript variants encoding two isoforms with different length have been identified. [provided by RefSeq, Aug 2010]


Synonyms: CEBPZ; CHOP; CHOP-10; CHOP10; GADD153

Protein Families: Druggable Genome, Transcription Factors

**Protein Pathways:** MAPK signaling pathway

# **Product images:**





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DDIT3 ([RC201301], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DDIT3. Positive lysates [LY418226] (100ug) and [LC418226] (20ug) can be purchased separately from OriGene.

Western blot analysis of HT29 cell lysate (35ug) by using anti-DDIT3 monoclonal antibody. Dilution: 1:500