

Product datasheet for TA802210S

DDIT3 Mouse Monoclonal Antibody [Clone ID: OTI3H7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3H7
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human DDIT3 (NP_004074) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	19 kDa
Gene Name:	DNA damage inducible transcript 3
Database Link:	<u>NP_004074</u> <u>Entrez Gene 13198 MouseEntrez Gene 29467 RatEntrez Gene 1649 Human</u> <u>P35638</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


DDIT3 Mouse Monoclonal Antibody [Clone ID: OTI3H7] – TA802210S

- Background:This gene encodes a member of the CCAAT/enhancer-binding protein (C/EBP) family of
transcription factors. The protein functions as a dominant-negative inhibitor by forming
heterodimers with other C/EBP members, such as C/EBP and LAP (liver activator protein), and
preventing their DNA binding activity. The protein is implicated in adipogenesis and
erythropoiesis, is activated by endoplasmic reticulum stress, and promotes apoptosis. Fusion
of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by
translocation generates chimeric proteins in myxoid liposarcomas or Ewing sarcoma.
Multiple alternatively spliced transcript variants encoding two isoforms with different length
have been identified. [provided by RefSeq, Aug 2010]
- Synonyms:CEBPZ; CHOP; CHOP-10; CHOP10; GADD153Protein Families:Druggable Genome, Transcription Factors


MAPK signaling pathway

Protein Pathways:

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DDIT3 (Cat# [RC201301], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DDIT3(Cat# [TA802210]). Positive lysates [LY418226] (100ug) and [LC418226] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue tissue using anti-DDIT3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA802210]) (1:500)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US




Immunohistochemical staining of paraffinembedded Human lung tissue within the normal limits using anti-DDIT3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA802210]) (1:500)

Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-DDIT3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA802210]) (1:500)

Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-DDIT3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA802210]) (1:500)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-DDIT3 mouse monoclonal antibody. (Heatinduced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA802210]) (1:500)

Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-DDIT3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA802210]) (1:500)

Immunohistochemical staining of paraffinembedded Human Gastric Carcinoma using anti-DDIT3 mouse monoclonal antibody. (Heatinduced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA802210]) (1:500)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US