

# Product datasheet for TA802377AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## LKB1 (STK11) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1F10]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1F10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-236 of human STK11

(NP\_000446) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 48.5 kDa

**Gene Name:** serine/threonine kinase 11

Database Link: NP 000446

Entrez Gene 20869 MouseEntrez Gene 314621 RatEntrez Gene 6794 Human

Q15831

**Background:** This gene, which encodes a member of the serine/threonine kinase family, regulates cell

polarity and functions as a tumor suppressor. Mutations in this gene have been associated with Peutz-Jeghers syndrome, an autosomal dominant disorder characterized by the growth of polyps in the gastrointestinal tract, pigmented macules on the skin and mouth, and other neoplasms. Alternate transcriptional splice variants of this gene have been observed but have

not been thoroughly characterized. [provided by RefSeq, Jul 2008]






**Synonyms:** hLKB1; LKB1; PJS

**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Adipocytokine signaling pathway, mTOR signaling pathway

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY STK11 ([RC202885], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-STK11. Positive lysates [LY424705] (100ug) and [LC424705] (20ug) can be purchased separately from OriGene.