

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA802569BM

PARN Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4D6]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI4D6
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-266 of human PARN (NP_002573) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	73.3 kDa
Gene Name:	poly(A)-specific ribonuclease
Database Link:	<u>NP_002573</u> <u>Entrez Gene 74108 MouseEntrez Gene 5073 Human</u> <u>O95453</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Serigene PARN Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4D6] – TA802569BM

Background:The protein encoded by this gene is a 3'-exoribonuclease, with similarity to the RNase D
family of 3'-exonucleases. It prefers poly(A) as the substrate, hence, efficiently degrades
poly(A) tails of mRNAs. Exonucleolytic degradation of the poly(A) tail is often the first step in
the decay of eukaryotic mRNAs. This protein is also involved in silencing of certain maternal
mRNAs during oocyte maturation and early embryonic development, as well as in nonsense-
mediated decay (NMD) of mRNAs that contain premature stop codons. Alternatively spliced
transcript variants encoding different isoforms have been found for this gene. [provided by
RefSeq, Aug 2008]

Synonyms:	DAN
Protein Families:	Transcription Factors
Protein Pathways:	RNA degradation

Product images:

170	-	
130	-	
100	-	_
70	-	-
55	-	
40	-	
35	-	
25	-	
15	-	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PARN ([RC207220], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PARN. Positive lysates [LY419233] (100ug) and [LC419233] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US