

Product datasheet for TA802590M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ornithine Carbamoyltransferase (OTC) Mouse Monoclonal Antibody [Clone ID: OTI2G6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2G6
Applications: IHC, WB
Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 33-354 of human OTC

(NP_000522)produced in SF9 cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 36.1 kDa

Gene Name: ornithine carbamoyltransferase

Database Link: NP 000522

Entrez Gene 18416 MouseEntrez Gene 25611 RatEntrez Gene 5009 Human

P00480

Background: This nuclear gene encodes a mitochondrial matrix enzyme. Missense, nonsense, and

frameshift mutations in this enzyme lead to ornithine transcarbamylase deficiency, which causes hyperammonemia. Since the gene for this enzyme maps close to that for Duchenne muscular dystrophy, it may play a role in that disease also. [provided by RefSeq, Jul 2008]

Synonyms: OCTD; OTCD






Ornithine Carbamoyltransferase (OTC) Mouse Monoclonal Antibody [Clone ID: OTI2G6] – TA802590M


Protein Families: Druggable Genome

Protein Pathways: Arginine and proline metabolism, Metabolic pathways

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY OTC ([RC214662], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-OTC.



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-OTC mouse monoclonal antibody. ([TA802590]) Dilution: 1:150. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.