

## **Product datasheet for TA802692S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Ornithine Carbamoyltransferase (OTC) Mouse Monoclonal Antibody [Clone ID: OTI8A9]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI8A9
Applications: IHC, WB
Recommended Dilution: WB 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 33-354 of human OTC

(NP\_000522)produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 36.1 kDa

**Gene Name:** ornithine carbamoyltransferase

Database Link: NP 000522

Entrez Gene 18416 MouseEntrez Gene 25611 RatEntrez Gene 5009 Human

P00480

**Background:** This nuclear gene encodes a mitochondrial matrix enzyme. Missense, nonsense, and

frameshift mutations in this enzyme lead to ornithine transcarbamylase deficiency, which causes hyperammonemia. Since the gene for this enzyme maps close to that for Duchenne muscular dystrophy, it may play a role in that disease also. [provided by RefSeq, Jul 2008]

Synonyms: OCTD; OTCD






# Ornithine Carbamoyltransferase (OTC) Mouse Monoclonal Antibody [Clone ID: OTI8A9] – TA802692S


**Protein Families:** Druggable Genome

**Protein Pathways:** Arginine and proline metabolism, Metabolic pathways

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY OTC ([RC214662], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-OTC.



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-OTC mouse monoclonal antibody. ([TA802692]) Dilution: 1:150